



**IV Iron Replacement Therapy for Iron Deficiency Anemia
in Pregnancy (2nd & 3rd Trimester)
- OUTPATIENT ORDERS**

General

ALLERGY: _____

Diagnosis (check all that apply):

- O99.012 Anemia complicating pregnancy, second trimester
- O99.013 Anemia complicating pregnancy, third trimester
- D50.9 Iron deficiency anemia, unspecified
- D50.8 Other Iron Deficiency Anemias

- Metric Measurements: Weight: _____ kilograms; Height: _____ centimeters.
- OK to start infusion on date _____ based on labs dated _____
- Number of weeks gestation at time of order _____

Laboratory

Ensure ordering provider to include/ attach a copy of current (within 30 days) lab results with this order set: CBC, Iron Panel (Iron, Transferrin, % TRF) and Ferritin

Nursing Orders

- Hypersensitivity / Anaphylaxis Management for Outpatient Infusions / Desensitization Order-Set
Note: epinephrine and methylprednisolone (Solu-medrol) are pregnancy category C
- Vital signs: baseline and every 30 minutes until discharge
- Hold patient 30 minutes post-infusion to observe for any signs and symptoms of hypersensitivity reactions

Hydration Orders

Sodium chloride 0.9% 500 mL IV x 1 bag at 500 mL/hour as needed to decrease vein discomfort

Pre-Medications

- Acetaminophen (Tylenol) 650 mg orally x 1 dose (Pregnancy Category C)
- DiphenhydrAMINE (Benadryl) 25 mg orally IV x 1 dose (Pregnancy category B)
- Other _____

Medication Orders

Iron Sucrose (Venofer)

- Iron sucrose 200 mg in sodium chloride 0.9% 100 mL IV over 30 minutes once a week x 5 doses
- Iron sucrose 200 mg in sodium chloride 0.9% 100 mL IV over 30 minutes twice a week x 5 doses
- Other _____

MD Print name _____

MD Contact #: _____

MD Signature _____

DATE ____/____/____ TIME ____:____

Patient Name:
Date of Birth: