



General

ALLERGY: _____

PRIMARY DIAGNOSIS: *** Must choose a primary from lists below ***

- D50.0 Iron deficiency anemia secondary to blood loss (chronic) D64.81 Anemia due to antineoplastic chemotherapy
 D50.8 Other Iron Deficiency Anemias D63.1 Anemia in chronic kidney disease

SECONDARY DIAGNOSIS:

- K90.9 Intestinal malabsorption, unspecified N18.9 Chronic kidney disease, unspecified
 T50.905A Unspecified adverse effect of unspecified drug, medicinal and biological substance
 T50.995A Other drug allergy

Metric Measurements: Weight: _____ kilograms; Height: _____ centimeters.

OK to start infusion on date _____ based on labs dated _____

Laboratory

Ensure ordering provider to include/ attach a copy of current (within 30 days old) lab results (baseline CBC, Iron Panel (Iron, Transferrin, % TRF, and Ferritin) with this order-set

Nursing Orders

IV ACCESS: Peripheral PICC Tunneled Catheter Implanted Port Other _____
 May Access Port-A-Cath May draw labs from Central Line

Hypersensitivity / Anaphylaxis Management for Outpatient Infusions / Desensitizations Order-Set

Vital signs: baseline and every 30 minutes until discharge

Hold patient 30 minutes post-infusion to observe for any signs and symptoms of hypersensitivity reactions

Hydration Orders

Sodium chloride 0.9% 500 mL IV x 1 bag at 500 mL/hour as needed to decrease vein discomfort

Pre-Medications

- Acetaminophen (Tylenol) 650 mg orally x 1 dose Famotidine (Pepcid) 20 mg IV x 1 dose
 DiphenhydrAMINE (Benadryl) 25 mg IV x 1 dose Hydrocortisone (Solu-cortef) 100 mg IV x 1 dose
or if not available methylprednisolone (Solu-medrol) 125 mg IV x 1 dose

Medication Orders

Ferric Derisomaltose (Monoferric)

Ferric Derisomaltose 1000mg in sodium chloride 0.9% 250 mL IV over 20 minutes x 1 dose. If Monoferric is not the preferred agent, may use (Venofer) Iron sucrose 300 mg in sodium chloride 0.9% 250 mL IV over 90 minutes every 2 days x 3 doses

If weight less than 50 kg, pharmacist to calculate dose at 20 mg/kg of body weight

Ferric Carboxymaltose (Injectafer)

Ferric carboxymaltose 750 mg in sodium chloride 0.9% 250 mL IV over 30 minutes x 1 dose
 Ferric carboxymaltose 750 mg in sodium chloride 0.9% 250 mL IV over 30 minutes every 7 days x 2 doses

If weight less than 50 kg, pharmacist to notify MD for dose calculation at 15 mg/kg of body weight

Ferumoxytol (Feraheme) for chronic kidney disease, unspecified (D63.1/N18.9) ONLY

Ferumoxytol 510 mg in sodium chloride 0.9% 100 mL IV over 30 minutes x 1 dose
 Ferumoxytol 510 mg in sodium chloride 0.9% 100 mL IV over 30 minutes every 7 days x 2 doses

Iron Sucrose (Venofer)

Iron sucrose 300 mg in sodium chloride 0.9% 250 mL IV over 90 minutes every 2 days x 3 doses

MD Print name _____ MD Contact #: _____

MD Signature _____ DATE ____/____/____

Reviewed: 02/2023

Patient Name:
Date of Birth: